

Table 10.3

Kaposi Sarcoma

Number of Cases

By SEER Location and Year of Diagnosis, 1975-2011

All Races, Males, Ages 55+

Year of Diagnosis

SEER Location	1975-1979	1980-1981	1982-1983	1984-1985	1986-1987	1988-1989	1990-1991	1992-1993	1994-1995	1996-1997	1998-1999	2000-2001	2002-2003	2004-2005	2006-2007	2008-2009	2010-2011
SEER 9 ^b	121	53	71	82	93	135	129	134	126	85	73	53	74	61	65	89	91
SEER 9 except SF-Oakland	91	39	49	54	49	81	69	65	65	53	47	37	39	42	41	63	69
San Francisco-Oakland SMSA ^c	30	14	22	28	44	54	60	69	61	32	26	16	35	19	24	26	22
San Francisco City/County	14	<4	8	20	31	35	38	44	42	17	10	7	18	10	9	12	11

Age-Adjusted Cancer Incidence Rates^a

By SEER Location and Year of Diagnosis, 1975-2011

All Races, Males, Ages 55+

Year of Diagnosis

SEER Location	1975-1979	1980-1981	1982-1983	1984-1985	1986-1987	1988-1989	1990-1991	1992-1993	1994-1995	1996-1997	1998-1999	2000-2001	2002-2003	2004-2005	2006-2007	2008-2009	2010-2011
SEER 9 ^b	1.9	1.9	2.4	2.5	2.5	3.6	3.6	3.3	3.0	2.1	1.7	1.3	1.5	1.2	1.2	1.6	1.5
SEER 9 except SF-Oakland	1.7	1.7	2.0	2.0	1.8	2.7	2.4	2.0	1.9	1.6	1.3	1.1	1.0	1.0	0.9	1.3	1.3
San Francisco-Oakland SMSA ^c	2.9	3.1	4.6	4.7	6.6	8.1	9.6	10.4	9.0	4.8	3.7	2.3	4.5	2.3	2.8	2.9	2.3
San Francisco City/County	4.7	-	5.6	12.9	20.5	23.5	26.4	30.4	29.8	11.5	6.7	4.4	10.8	5.7	4.8	6.3	5.2

^a Rates are per 100,000 males aged 55+ and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).^b The SEER 9 areas are San Francisco-Oakland SMSA, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah and Atlanta.^c The San Francisco-Oakland SMSA represents 15% of all persons in the SEER areas. For this age group, all SEER areas may not be generalizable to the United States for Kaposi Sarcoma.

- Statistic not shown. Rate based on less than 4 cases for the time interval.